

**Arizona Bone and Joint Specialists**

5620 E. Bell Rd, Scottsdale, Arizona 85254 602.493.9361

13640 N. 7th Street, Phoenix, Arizona 85022 602.863.1962

**Foot/Leg**

## Plantar fasciitis

When your first few steps out of bed in the morning cause severe pain in the heel of your foot, you may have plantar fasciitis (fashee-EYE-tiss). It's an overuse injury affecting the sole or flexor surface (plantar) of the foot. A diagnosis of plantar fasciitis means you have inflamed the tough, fibrous band of tissue (fascia) connecting your heel bone to the base of your toes.

You're more likely to get the condition if you're a woman, if you're overweight, or if you have a job that requires a lot of walking or standing on hard surfaces. You're also at risk if you walk or run for exercise, especially if you have tight calf muscles that limit how far you can flex your ankles. People with very flat feet or very high arches are also more prone to plantar fasciitis.


The condition starts gradually with mild pain at the heel bone often referred to as a stone bruise. You're more likely to feel it after (not during) exercise. The pain classically occurs again after arising from a midday lunch break.

If you don't treat plantar fasciitis, it may become a chronic condition. You may not be able to keep up your level of activity and you may also develop symptoms of foot, knee, hip and back problems because of the way plantar fasciitis changes the way you walk.

### Treatments

Rest is the first treatment for plantar fasciitis. Try to keep weight off your foot until the inflammation goes away. You can also apply ice to the sore area for 20 minutes three or four times a day to relieve your symptoms. Often a doctor will prescribe nonsteroidal anti-inflammatory medication such as ibuprofen. A program of home exercises to stretch your Achilles tendon and plantar fascia are the mainstay of treating the condition and lessening the chance of recurrence.

In one exercise, you lean forward against a wall with one knee straight and heel on the ground. Your other knee is bent. Your heel




cord and foot arch stretch as you lean. Hold for 10 seconds, relax and straighten up. Repeat 20 times for each sore heel.

In the second exercise, you lean forward onto a countertop, spreading your feet apart with one foot in front of the other. Flex your knees and squat down, keeping your heels on the ground as long as possible. Your heel cords and foot arches will stretch as the heels come up in the stretch. Hold for 10 seconds, relax and straighten up. Repeat 20 times.



About 90 percent of people with plantar fasciitis improve significantly after two months of initial treatment. You may be advised to use shoes with shock-absorbing soles or fitted with a standard orthotic device like a rubber heel pad. Your foot may be taped into a specific position.

If your plantar fasciitis continues after a few months of conservative treatment, your doctor may inject your heel with steroidal anti-inflammatory medications (corticosteroid). If you still have symptoms, you may need to wear a walking cast for 2-3 weeks or positional splint when you sleep. In a few cases, you might need surgery to release your ligament.



April 2001



Codeveloped by the [American Orthopaedic Foot and Ankle Society](#)

For More Information:

[The Foot and Ankle](#)

[Heel Pain](#)

[Common Foot Problems](#)

### Instructions to Patient:

---

[Main Menu](#) [Foot](#)